arXiv:1603.00334v1 [math.AC] 1 Mar 2016

.

FINITE F-TYPE AND F-ABUNDANT MODULES

HAILONG DAO AND TONY SE

Dedicated to Professor Kei-ichi Watanabe on the occasion of his seventieth birthday

Abstract. In this note we introduce and study basic properties of two types of modules over a commutative noetherian ring R of positive prime characteristic. The first is the category of modules of finite F-type. These objects include reflexive ideals representing torsion elements in the divisor class group of R. The second class is what we call F-abundant modules. These include, for example, the ring R itself and the canonical module when R has positive splitting dimension. We prove various facts about these two categories and how they are related, for example that $\mathrm{Hom}_R(M,N)$ is maximal Cohen-Macaulay when M is of finite F-type and N is F-abundant, plus some extra (but necessary) conditions. Our methods allow us to extend previous results by Patakfalvi-Schwede, Yao and Watanabe. They also afford a deeper understanding of these objects, including complete classifications in many cases of interest, such as complete intersections and invariant subrings.

1. Introduction

Let $(R,\mathfrak{m},k)$ be a reduced F-finite local ring of dimension d and prime characteristic $p>0$. Let $\alpha(R)=\log_p[k:k^p]$. All modules are in $\mathrm{mod}(R)$, the category of finitely generated R-modules.

In this paper we prove some connections between two types of objects defined using the Frobenius endomorphism $R\to R$ with $r\mapsto r^p$.

These objects generalize some well-studied concepts. Let us start with the definitions. Fix an R-module M. For $e\in\mathbb{Z}_{\geqslant 0}$, let ${}^eM$ denote the abelian group M viewed as an R-module via the eth iteration of the Frobenius map. We let $F^e\colon \mathrm{mod}(R)\to\mathrm{mod}(R)$ denote the eth Peskine-Szpiro functor given by

$$F^e(M):=M\otimes_R {}^eR.$$

Given $S\subseteq\mathrm{mod}(R)$, we use $\mathrm{add}_R(S)$ to denote the additive subcategory of $\mathrm{mod}(R)$ generated by S.

Definition 1.1. (1) Let M be an R-module such that $\mathrm{Supp}(M)=\mathrm{Spec}\,R$ and is locally free in codimension 1. We let $M(e)=F_R^e(M)^{**}$, the reflexive hull of $F_R^e(M)$, viewed as an R-module by identifying ${}^eR$ with R. We say that M is of finite F-type if $\{M(e)\}_{e\geqslant 0}\subseteq \mathrm{add}_R(X)$ for some R-module X (see Lemma 4.3). We let $\mathcal{FT}(R)$ denote the category of R-modules of finite F-type.

Date: March 2, 2016.

1991 *Mathematics Subject Classification.* Primary: 13A35; Secondary:13D07, 13H10.

Key words and phrases. Frobenius endomorphism, divisor class group, F-regularity.

The first author is partially supported by NSF grant 1104017.

(2) Let N, L be R-modules. Let b_e be maximum such that ${}^eN = L^{\oplus b_e} \oplus N_e$ for some N_e. We say that (N, L) is an abundant pair if $\liminf_{e\to\infty} p^{e\alpha(R)}/b_e = 0$.
(3) Let L be an R-module. We say that L is an F-abundant module if (N, L) is an abundant pair for some N.

Examples of modules of finite F-type include torsion elements of the divisor class group of a normal domain R (without any assumption about the order of the element), finite integral extensions that are étale in codimension one (see section 4), or F-periodic vector bundles on the punctured spectrum of R (and of the corresponding projective variety X when R is a local cone of some embedding of X; see section 6). For F-abundant modules, R has positive splitting dimension if and only if (R, R) is an abundant pair (see section 5). A good source of examples in both cases are the rings of invariants of a finite group, see Example 5.3.

Our main technical results (collected in section 3) say roughly that under various extra conditions, if M is of finite F-type and N is F-abundant, then $\operatorname{Hom}_R(M, N)$ is maximal Cohen-Macaulay. We shall give plenty of examples to show that the technical conditions are necessary. Our approach yields a strengthening of some well-known results as well as many new ones:

(1) An extension of results by Patakfalvi-Schwede ([18, Theorem 3.1]) and Watanabe ([21, Corollary 2.9]) on depth of divisor classes associated to F-regular singularities. See Theorem 6.3.
(2) Under certain conditions, a strong generalization of Yao's result ([22, Lemma 2.2]) on Cohen-Macaulayness of F-contributors. See Theorems 3.8 and 3.10.
(3) A complete classification of the categories of finite F-type R-modules and abundant F-modules in many cases of interest, such as complete intersections and invariant subrings (Corollary 4.13, Theorem 4.14 and 5).

2. Notation and preliminary results

Definition 2.1. ([3, Definition 2.4] and [5, Definition 4.5]) Let $a_e = a_e(R)$ be maximum such that ${}^eR = R^{\oplus a_e} \oplus R_e$ for some R_e. The largest integer k such that

$$\lim_{e\to\infty} \frac{a_e}{p^{e(k+\alpha(R))}} > 0$$

is called the F-splitting dimension of R, and is denoted by $\operatorname{sdim} R$.

Notation 2.2. (a) We use (S_k) to denote Serre's criteria for $k \geqslant 0$.
(b) We use $\mathrm{H}^i_{\mathfrak{m}}(M)$ to denote the ith local cohomology module of an R-module M supported on $\mathfrak{m}$.
(c) For an R-module M, we let M^* denote $\operatorname{Hom}_R(M, R)$.
(d) Suppose that R has a canonical module ω_R. Then for an R-module M, we let $M^\vee$ denote $\operatorname{Hom}_R(M, \omega_R)$.
(e) For $S \subseteq R$, we let ${}^eS$ to denote S viewed as a subset of ${}^eR$. Then for $P \in \operatorname{Spec} R$, ${}^eR_{e^P}$ is an R_P-module, and ${}^eR_{e^P} = {}^e(R_P)$.
(f) For an ideal $I \subseteq R$, an integer $e \geqslant 0$ and $q = p^e$, we let $I^{[q]} = \{x^q \mid x \in I\}$, the qth Frobenius power of I.

(g) We use $\mathrm{CM}(R)$ to denote the subcategory of $\mathrm{mod}(R)$ consisting of all C-M R-modules and $\mathrm{Ref}(R)$ the subcategory of all reflexive modules.

Remark 2.3. We would like to remind the reader of another characterization of $\operatorname{sdim} R$. We consider the splitting prime $\mathcal{P}(R)$ of R, as defined in [3, Definition 3.2]. By [3, Theorem 3.3 and Corollary 3.4], $\mathcal{P}(R)$ is a prime ideal if $\operatorname{sdim}(R) \neq -\infty$ or the unit ideal otherwise. Corollary 4.3 of [5] shows that $\operatorname{sdim} R = \dim(R/\mathcal{P}(R))$ when $\operatorname{sdim}(R) \neq -\infty$.

Proposition 2.4. *Assume the following for R:*
(1) R is equidimensional;
(2) R_P is C-M for all $P \in \operatorname{Spec} R \setminus \{\mathfrak{m}\}$; and
(3) $\operatorname{sdim} R > 0$.
Then R is C-M.

Proof. Since R is F-finite, it is a homomorphic image of a Gorenstein ring by [11, Remark before Lemma A.2]. By (1) and (2), for $0 \leqslant i < d$ we have that $\mathrm{H}^i_{\mathfrak{m}}(R)$ has finite length by [1, Proposition 21.24]; see also [2, Theorem 9.5.2 (Grothendieck's Finiteness Theorem)]. So

$$\begin{aligned} p^{e\alpha(R)}\lambda_R(\mathrm{H}^i_{\mathfrak{m}}(R)) &= \lambda_R({}^e\mathrm{H}^i_{\mathfrak{m}}(R)) \\ &= \lambda_R(\mathrm{H}^i_{\mathfrak{m}}({}^eR)) \\ &= \lambda_R(\mathrm{H}^i_{\mathfrak{m}}(R^{\oplus a_e} \oplus M_e)) \\ &= a^e\lambda_R(\mathrm{H}^i_{\mathfrak{m}}(R)) + \lambda_R(\mathrm{H}^i_{\mathfrak{m}}(M_e)) \\ \frac{1}{p^e}\lambda_R(\mathrm{H}^i_{\mathfrak{m}}(R)) &= \frac{a_e}{p^{e(1+\alpha(R))}}\lambda_R(\mathrm{H}^i_{\mathfrak{m}}(R)) + \frac{1}{p^{e(1+\alpha(R))}}\lambda_R(\mathrm{H}^i_{\mathfrak{m}}(M_e)) \\ 0 = \lim_{e\to\infty}\frac{1}{p^e}\lambda_R(\mathrm{H}^i_{\mathfrak{m}}(R)) &\geqslant \lim_{e\to\infty}\frac{a_e}{p^{e(1+\alpha(R))}}\lambda_R(\mathrm{H}^i_{\mathfrak{m}}(R)) \end{aligned}$$

Since $\operatorname{sdim} R \geqslant 1$, we have $\lambda_R(\mathrm{H}^i_{\mathfrak{m}}(R)) = 0$ and hence $\mathrm{H}^i_{\mathfrak{m}}(R) = 0$ for $0 \leqslant i < d$. Then by [1, Theorem 10.36], R is C-M. □

Corollary 2.5. *If R is F-split, is C-M on $\operatorname{Spec} R \setminus \{\mathfrak{m}\}$ (e.g. R is an isolated singularity) but not C-M, then $\operatorname{sdim} R = 0$.* □

Example 2.6. *Let X be an ordinary Abelian variety of dimension at least two and R be the coordinate ring of an embedding of X with respect to some polarization. It is well-known, as in* [15, 19]*, that R is F-split but not C-M. So by Corollary 2.5, $\operatorname{sdim} R = 0$.*

Lemma 2.7. *Let M, N be R-modules such that ${}^eM = N^{b_e} \oplus M_e$ and $\liminf_{e\to\infty} \frac{b_e}{p^{e(k+\alpha(R))}} > 0$. Then $\operatorname{depth} N \geqslant k$. In particular, if $k = \dim(M)$, then N is C-M.*

Proof. We use the same proof as in [22, Lemma 2.2]. For $0 \leqslant i < k$, we have

$$0 = \lim_{e\to\infty}\frac{\lambda_R(\mathrm{H}^i_R(\underline{x}^{p^e}, M))}{p^{ek}} = \lim_{e\to\infty}\frac{\lambda_R(\mathrm{H}^i_R(\underline{x}, {}^eM))}{p^{e(k+\alpha(R))}} \geqslant \liminf_{e\to\infty}\frac{b_e}{p^{e(k+\alpha(R))}}\lambda_R(\mathrm{H}^i_R(\underline{x}, N))$$

as in Proposition 2.4. So $\mathrm{H}^i_R(\underline{x}, N) = 0$ for all $0 \leqslant i < k$, and hence $\operatorname{depth} N \geqslant k$. □

Remark 2.8. It is already known that $\operatorname{sdim} R = d \Rightarrow R$ is strongly F-regular $\Rightarrow R$ is C-M.

Remark 2.9. Let ${}^eM = R^{b_e} \oplus M_e$ with b_e the largest possible. The largest integer k such that

$$\liminf_{e\to\infty} \frac{b_e}{p^{e(k+\alpha(R))}} > 0$$

was defined in [3, Definition 5.4] to be $\operatorname{sdim}(M)$, the s-dimension of M.

Definition 2.10. We will let (sdim_n) to denote the statement: $\operatorname{sdim} R_P > 0$ for all $P \in \operatorname{Spec} R$ such that $\operatorname{ht} P \geqslant n$.

Lemma 2.11. *If R satisfies (sdim_n), then $\operatorname{ht} \mathcal{P}(R) < n$.*

Proof. Suppose that R satsifies (sdim_n). By [3, Proposition 3.6], we have $\mathcal{P}(R_{\mathcal{P}(R)}) = \mathcal{P}(R)R_{\mathcal{P}(R)}$. Since R is F-finite and reduced, so is $R_{\mathcal{P}(R)}$. By [3, Corollary 3.4], since $\mathcal{P}(R_{\mathcal{P}(R)})$ is the maximal ideal of $R_{\mathcal{P}(R)}$, we have $\operatorname{sdim} R_{\mathcal{P}(R)} = 0$, so $\operatorname{ht} \mathcal{P}(R) < n$. □

3. Main technical results

Lemma 3.1. *Let $f\colon M \to N$ be a homomorphism of R-modules, where M is (S_2) and N is (S_1). Suppose that f is an isomorphism in codimension 1. Then f is an isomorphism.*

Remark 3.2. In many of the results below, we can replace the assumption of a module being locally free in codimension 1 by R being quasinormal.

Corollary 3.3. *Let L, N be R-modules such that L is free in codimension 1 and N is (S_2). Then*

$$\operatorname{Hom}_R(L, N) \cong \operatorname{Hom}_R(L^{**}, N)$$

Proof. Let $f\colon L \to L^{**}$ be the canonical map and $\overline{f}\colon \operatorname{Hom}_R(L^{**}, N) \to \operatorname{Hom}_R(L, N)$ the induced map. Then $\overline{f}$ is an isomorphism in codimension 1. By Lemma 3.1, it suffices to show that $\operatorname{Hom}_R(M, N)$ is (S_2) for every R-module M. Given M, let $F_1 \to F_0 \to M \to 0$ be a finite presentation of M. Applying $\operatorname{Hom}_R(-, N)$ gives

$$0 \to \operatorname{Hom}_R(M, N) \to \operatorname{Hom}_R(F_0, N) \to \operatorname{Hom}_R(F_1, N)$$

Since N is (S_2), so are $\operatorname{Hom}_R(F_i, N)$ for $i = 1, 2$ and $\operatorname{Hom}_R(M, N)$, as required. □

Corollary 3.4. *Suppose that R is C-M with a canonical module ω_R. Suppose that M is (S_2) and $M^\vee$ is MCM. Then M is MCM.*

Proof. Consider the natural map $f\colon M \to M^{\vee\vee}$. In codimension 1, M is MCM since it is (S_2), so f is an isomorphism. Since $M^\vee$ is MCM, so is $M^{\vee\vee}$. Hence f is an isomorphism by Lemma 3.1. □

Remark 3.5. Let M be an R-module, $f\colon R \to S$ be a ring homomorphism, U be a multiplicative subset of S and $T = f^{-1}(U)$. Then we have $(M \otimes_R S)_U = M \otimes_R S \otimes_S S_U = M \otimes_R S_U = M \otimes_R R_T \otimes_{R_T} S_U = M_T \otimes_{R_T} S_U$.

Lemma 3.6. *Let $R \xrightarrow{f_1} R_1 \xrightarrow{f_2} R_2$ be a sequence of ring homomorphisms where R_2 is (S_2). Let $f = f_2 \circ f_1$. Let M be an R-module. Suppose that M_P is free for every $P \in \operatorname{Spec} R$ of height 1 and for every $P = f^{-1}(Q)$ such that $Q \in \operatorname{Spec} R_2$ of height 1. Let $M_i = (M \otimes_R R_i)^{**}$ (over R_i) for $i = 1, 2$. Then $(M_1 \otimes_{R_1} R_2)^{**} \cong M_2$.*

Proof. Let $N = M \otimes_R R_1$. The natural map $N \to M_1 = N^{**}$ gives rise to a map $g\colon M \otimes_R R_2 = M \otimes_R R_1 \otimes_{R_1} R_2 = N \otimes_{R_1} R_2 \to N^{**} \otimes_{R_1} R_2 = M_1 \otimes_{R_1} R_2$. Let $Q \in \operatorname{Spec} R_2$, $P_1 = (f_2)^{-1}(Q)$ and $P = f^{-1}(Q)$. Then $(M \otimes_R R_2)_Q = M_P \otimes_{R_P} (R_2)_Q$ and $(M_1 \otimes_{R_1} R_2)_Q = (M \otimes_R R_1)^{**}_{P_1} \otimes_{(R_1)_{P_1}} (R_2)_Q = (M_P \otimes_{R_P} (R_1)_{P_1})^{**} \otimes_{(R_1)_{P_1}} (R_2)_Q$, so g is an isomorphism in codimension 1. Applying ** gives the map $M_2 = (M \otimes_R R_2)^{**} \to (M_1 \otimes_{R_1} R_2)^{**}$, which is an isomorphism by Lemma 3.1 since R_2 is (S_2). □

Corollary 3.7. *("Index shifting") Let R be (S_2) and M be as in Definition 1.1. Let e, f be nonnegative integers. Then $[M(e)](f) \cong M(e+f)$.*

Proof. Let $R_1 = {}^eR$ and $R_2 = {}^{e+f}R$ as in Lemma 3.6. Then we have $M(e) = M_1$, and so $[M(e)](f) = (M_1 \otimes_{R_1} R_2)^{**} \cong M_2 = M(e+f)$. □

Theorem 3.8. *Suppose that R is (S_2) and equidimensional. Let M, N be R-modules such that $M \in \mathcal{FT}(R)$ and N is (S_2). Assume that (N, L) is an abundant pair and that $N_P \in \operatorname{add} L_P$ for all $P \in \operatorname{Spec} R$ such that $3 \leqslant \operatorname{ht}(P) < d$. Assume further that for every $P \in \operatorname{Spec} R$ such that $3 \leqslant \operatorname{ht}(P) < d$ and $e \geqslant 0$, $(\operatorname{Hom}_R(M(e), L))_P$ is MCM. Then $\operatorname{Hom}_R(M(e), L)$ is MCM for all $e \geqslant 0$.*

Proof. If $d \leqslant 2$, then $\operatorname{Hom}_R(M(e), L)$ is MCM since L is (S_2). So we may assume that $d \geqslant 3$. By assumption, for $P \in \operatorname{Spec} R \setminus \{\mathfrak{m}\}$, we have $N_P \in \operatorname{add} L_P$ for $\operatorname{ht}(P) \geqslant 3$, so $(\operatorname{Hom}_R(M(e), N))_P$ is MCM for $e \geqslant 0$. So for $0 \leqslant i < d$ and $e \geqslant 0$, $\operatorname{H}^i_{\mathfrak{m}}(\operatorname{Hom}_R(M(e), N))$ has finite length — see Proposition 2.4. By Corollary 3.3, we have

$$\begin{aligned}
\operatorname{Hom}_{{}^eR}({}_{{}^eR}[F^e_R(M)^{**}], {}_{{}^eR}N) &= \operatorname{Hom}_{{}^eR}({}_{{}^eR}[(M \otimes_R {}^eR)^{**}], {}_{{}^eR}N) \\
&= \operatorname{Hom}_{{}^eR}({}_{{}^eR}[M \otimes_R {}^eR], {}_{{}^eR}N) \\
&= \operatorname{Hom}_R(M, \operatorname{Hom}_{{}^eR}({}^eR, {}_{{}^eR}N)) \\
&= \operatorname{Hom}_R(M, {}_{{}^eR}N) \\
&= \operatorname{Hom}_R(M, {}^eN) \\
&= \operatorname{Hom}_R(M, L^{\oplus b_e} \oplus N_e) \\
&= \operatorname{Hom}_R(M, L)^{\oplus b_e} \oplus \operatorname{Hom}_R(M, N_e)
\end{aligned}$$

Apply $\operatorname{H}^i_{\mathfrak{m}}$ for $0 \leqslant i < d$ to get

$$\begin{aligned}
b_e \lambda_R(\operatorname{H}^i_{\mathfrak{m}}(\operatorname{Hom}_R(M, L))) &\leqslant \lambda_R(\operatorname{H}^i_{\mathfrak{m}}(\operatorname{Hom}_{{}^eR}({}_{{}^eR}[F^e_R(M)^{**}], {}_{{}^eR}N))) \\
&= \lambda_R(\operatorname{H}^i_{\mathfrak{m}({}^eR)}(\operatorname{Hom}_{{}^eR}({}_{{}^eR}[F^e_R(M)^{**}], {}_{{}^eR}N))) \quad \text{(base change)} \\
&= \lambda_R(\operatorname{H}^i_{{}^e\mathfrak{m}}(\operatorname{Hom}_{{}^eR}({}_{{}^eR}[F^e_R(M)^{**}], {}_{{}^eR}N))) \\
&= \lambda_R({}^e\operatorname{H}^i_{\mathfrak{m}}(\operatorname{Hom}_R(M(e), {}_RN))) \\
&= p^{e\alpha(R)} \lambda_R(\operatorname{H}^i_{\mathfrak{m}}(\operatorname{Hom}_R(M(e), {}_RN))) \\
\lambda_R(\operatorname{H}^i_{\mathfrak{m}}(\operatorname{Hom}_R(M, L))) &\leqslant \frac{p^{e\alpha(R)}}{b_e} \max\{\lambda_R(\operatorname{H}^i_{\mathfrak{m}}(\operatorname{Hom}_R(M(e), {}_RN)))\}
\end{aligned}$$

Taking lim inf shows that $\operatorname{H}^i_{\mathfrak{m}}(\operatorname{Hom}_R(M, L)) = 0$ for $0 \leqslant i < d$, so $\operatorname{Hom}_R(M, L)$ is MCM. By Corollary 3.7, we have $[M(e)](f) \cong M(e+f)$. So we may replace M by $M(e)$ to conclude that $\operatorname{Hom}_R(M(e), L)$ is MCM. □

Remark 3.9. By Lemma 4.3, we only need to check that $(\operatorname{Hom}_R(M_i, L))_P$ is MCM for $3 \leqslant \operatorname{ht}(P) < d$ for the finitely many indecomposable modules M_i that appear among $\{M(e)\}_{e\geqslant 0}$.

Theorem 3.10. *Let R be as in Theorem 3.8. Let M, N be R-modules such that $M \in \mathcal{FT}(R)$ and N is (S_2). Assume that for every $P \in \operatorname{Spec} R$ such that $\operatorname{ht}(P) \geqslant 3$, (N_P, L_P) is an abundant pair. Assume further that for every $P \in \operatorname{Spec} R$ such that $3 \leqslant \operatorname{ht}(P) < d$, we have $N_P \in \operatorname{add} L_P$. Then $\operatorname{Hom}_R(M(e), L)$ is MCM for all $e \geqslant 0$.*

Proof. First, for every $P \in \operatorname{Spec} R$, we have the following.

$$\begin{aligned}
{}^{e(R_P)}[F^e_{R_P}(M_P)^{**}] &= \operatorname{Hom}_{e(R_P)}\left(\operatorname{Hom}_{e(R_P)}\left(M_P \otimes_{e(R_P)} {}^{e}(R_P), {}^{e}(R_P)\right), {}^{e}(R_P)\right)\\
&= \operatorname{Hom}_{e(R_P)}\left(\operatorname{Hom}_{e(R_P)}\left(M \otimes_R R_P \otimes_{e(R_P)} {}^{e}(R_P), {}^{e}(R_P)\right), {}^{e}(R_P)\right)\\
&= \operatorname{Hom}_{e(R_P)}\left(\operatorname{Hom}_{e(R_P)}\left(M \otimes_R {}^{e}(R_P), {}^{e}(R_P)\right), {}^{e}(R_P)\right)\\
&= \operatorname{Hom}_{e(R_P)}\left(\operatorname{Hom}_{{}^{e}R\,{}^{e}P}\left(M \otimes_R {}^{e}R \otimes_{{}^{e}R\,{}^{e}P} {}^{e}R_{{}^{e}P}, {}^{e}R_{{}^{e}P}\right), {}^{e}(R_P)\right)\\
&= \operatorname{Hom}_{e(R_P)}\left(\operatorname{Hom}_{{}^{e}R}\left(M \otimes_R {}^{e}R, {}^{e}R\right)_{{}^{e}P}, {}^{e}(R_P)\right)\\
&= \left(\operatorname{Hom}_{{}^{e}R}\left(\operatorname{Hom}_{{}^{e}R}\left(M \otimes_R {}^{e}R, {}^{e}R\right), {}^{e}R\right)\right)_{{}^{e}P}\\
&= (F^e_R(M)^{**})_P
\end{aligned}$$

So we can prove by induction on d that $\operatorname{Hom}_R(M(e), L)$ is MCM for all $e \geqslant 0$. We may assume that $d \geqslant 3$. Let $P \in \operatorname{Spec} R$ such that $3 \leqslant \operatorname{ht}(P) < d$. By induction, $(\operatorname{Hom}_R(M(e), L))_P$ is MCM. So by Theorem 3.8, $\operatorname{Hom}_R(M(e), L)$ is MCM. □

Corollary 3.11. *Suppose that R is C-M and $M \in \mathcal{FT}(R)$ is (S_2). Assume that:*
(a) either $\operatorname{sdim} R > 0$ and $M(e)_P$ is MCM for every $P \in \operatorname{Spec} R$ such that $3 \leqslant \operatorname{ht}(P) < d$ and $e \geqslant 0$; or
(b) R is (sdim_3).
Then M is MCM.

Proof. Since R is a homomorphic image of a Gorenstein ring, it has a canonical module ω_R by [2, 12.1.3(iii)]. Since R is C-M, it is equidimensional, and ω_R is reflexive and hence (S_2). We will show that $\operatorname{Hom}_R(M, \omega_R) = M^\vee$ is MCM, so that by Corollary 3.4, M is MCM. We may assume that $d > 2$. Since $\operatorname{sdim} R > 0$, (R, R) is an abundant pair by Example 5.1. Then (ω_R, ω_R) is also an abundant pair, since $\omega_{{}^{e}R} \cong \operatorname{Hom}_R({}^{e}R, \omega_R) = \operatorname{Hom}_R(R^{\oplus a_e} \oplus R_e, \omega_R) = \omega_R^{\oplus a_e} \oplus \operatorname{Hom}_R(R_e, \omega_R)$. If (a) holds, then $(\operatorname{Hom}_R(M(e), \omega_R))_P$ is MCM for every $P \in \operatorname{Spec} R$ such that $3 \leqslant \operatorname{ht}(P) < d$ and $e \geqslant 0$. By Theorem 3.8, $\operatorname{Hom}_R(M, \omega_R) = M^\vee$ is MCM. If (b) holds, then $(\omega_{R_P}, \omega_{R_P})$ is an abundant pair for $P \in \operatorname{Spec} R$ such that $\operatorname{ht}(P) \geqslant 3$, so $\operatorname{Hom}_R(M, \omega_R)$ is MCM by Theorem 3.10. □

Corollary 3.12. *Suppose that R is strongly F-regular and $M \in \mathcal{FT}(R)$ is (S_2). Then M is MCM. In particular, if I is a reflexive ideal such that $[I]$ is torsion in $\operatorname{Cl}(R)$, then I is MCM.*

Proof. Since R is strongly F-regular, it is (sdim_3). Corollary 3.11 shows that M is MCM. By Example 4.1, we have $I \in \mathcal{FT}(R)$. □

Remark 3.13. Corollary 3.12 generalizes [18, Corollary 3.3].

4. The category of finite F-type modules

In this section we study the category of finite F-type in more detail. We completely classify this category when R is a quotient singularity with a finite group whose order is coprime to p (Corollary 4.12), or when R is a complete intersection that is regular in codimension 2 (Theorem 4.14).

Example 4.1. *The following are R-modules M of finite F-type.*
(a) ([21, paragraph 2.3]) R is a normal domain and $M = I$, where I is a divisorial ideal. Then $M(e) \cong I^{(e)}$, so $M \in \mathcal{FT}(R)$ iff $[I]$ is torsion in $\mathrm{Cl}(R)$.
(a') M is a free R-module.
(b) ([21, Theorem 2.7]) Let $R \to S$ be a finite homomorphism of normal domains which is étale in codimension 1 and $M = S$. From the natural map $S \otimes_R {}^eR \to {}^eS$ we get $(S \otimes_R {}^eR)^{**} \cong S^{**}$, i.e. $M(e) \cong S$.
(c) See also Lemma 6.2.

Lemma 4.2. *$\mathcal{FT}(R)$ is closed under direct sums and direct summands.*

Proof. Obvious. □

Lemma 4.3. *Let $S \subseteq \operatorname{mod} R$. Then $\operatorname{add}_R(S)$ has finitely many indecomposable objects iff $S \subseteq \operatorname{add}_R(X)$ for some R-module X. Hence for an R-module M, $M \in \mathcal{FT}(R)$ if and only if only finitely many indecomposable direct summands appear among $\{M(e)\}_{e \geqslant 0}$.*

Proof. For the "only if" part, take X to be the direct sum of the indecomposable objects of $\operatorname{add}_R(S)$. For the "if" part, we consider the endomorphism ring $E = \operatorname{End}_R(X)$. Consider the category $\operatorname{Proj} E \subseteq \operatorname{mod}(E)$ of left projective modules over E. Then $F\colon \operatorname{add}_R(X) \to \operatorname{Proj} E$ given by $F(L) = \operatorname{Hom}_R(L, X)$ for $L \in \operatorname{add}_R(X)$ is an equivalence of categories. Since E is finitely generated over the local ring R, it is semilocal. By [8, Theorem 9], $\operatorname{Proj} E$ has only finitely many isomorphism classes of indecomposable objects, and hence so does $\operatorname{add}_R(X)$. □

Corollary 4.4. *Let R, M be as in Corollary 3.7. Then $M \in \mathcal{FT}(R)$ iff there are $e \geqslant 0$ and $f > 0$ such that $M(e) \cong M(e + f)$.*

Proof. "If": We note that $M(e + g) \cong [M(e)](g) \cong [M(e + f)](g) \cong M(e + f + g)$ by Corollary 3.7, so there are only finitely many isomorphism classes of $\{M(e)\}_{e \geqslant 0}$.
"Only if": Let X be an R-module such that $\{M(e)\}_{e \geqslant 0} \subseteq \operatorname{add}_R(X)$. Let S be the set of indecomposable direct summands of $\{M(e)\}_{e \geqslant 0}$. Then $S \subseteq \operatorname{add}_R(X)$, and S is finite by Lemma 4.3. By index shifting, it suffices to prove that $N(e) \cong N(e + f)$ for some $e \geqslant 0$ and $f > 0$ for all $N \in S$. By assumption, R is local, so we can prove the claim by induction on the minimum number of generators of N. Suppose first that $N(e)$ is indecomposable for all e. Then the claim holds since S is finite. So suppose that $N(e_0)$ is not indecomposable for some e_0 and has indecomposable direct summands $N_i \in S$. By induction, the claim holds for the N_i, and hence for N by index shifting. □

Proposition 4.5. *Let R be (S_2). Let $\mathcal{FT} = \mathcal{FT}(R)$. Then:*
(a) if $M \in \mathcal{FT}$, then $M^{**} \in \mathcal{FT}$.
(b) if $M, N \in \mathcal{FT}$, then $M \otimes N \in \mathcal{FT}$.

Proof. (a) By Corollary 3.7, we have $[M(0)](e) \cong M(e)$, that is, $(M^{**})(e) \cong M(e)$.
(b) For $P \in \operatorname{Spec} R$, we have $(M \otimes_R N)_P = M_P \otimes_{R_P} N_P$, so $M \otimes N$ is locally free in codimension 1. Next, we have

$$\begin{aligned} M \otimes_R N \otimes_R {}^eR &= M \otimes_R N \otimes_R {}^eR \otimes_{{}^eR} {}^eR \\ &= M \otimes_R {}^eR \otimes_{{}^eR} {}^eR \otimes_R N \\ &= (M \otimes_R {}^eR) \otimes_{{}^eR} (N \otimes_R {}^eR) \\ (M \otimes_R N \otimes_R {}^eR)^{**} &= [(M \otimes_R {}^eR) \otimes_{{}^eR} (N \otimes_R {}^eR)]^{**} \end{aligned}$$

The natural maps $M \otimes_R {}^eR \to (M \otimes_R {}^eR)^{**}$ and $N \otimes_R {}^eR \to (N \otimes_R {}^eR)^{**}$ give rise to the map $f\colon [(M \otimes_R {}^eR) \otimes_{{}^eR} (N \otimes_R {}^eR)]^{**} \to [(M \otimes_R {}^eR)^{**} \otimes_{{}^eR} (N \otimes_R {}^eR)^{**}]^{**}$. Since R is (S_2), Lemma 3.1 shows that f is an isomorphism. So we have an isomorphism

$$(M \otimes_R N)(e) \cong (M(e) \otimes_R N(e))^{**}$$

By Lemma 4.3, only finitely many indecomposable direct summands $\{K_i\}_{i=1}^m$ appear in $\{M(e)\}_{e \geqslant 0}$ and $\{L_j\}_{j=1}^n$ in $\{N(e)\}_{e \geqslant 0}$. Let $X = \sum_{i,j} (K_i \otimes_R L_j)^{**}$. Then $\{(M \otimes_R N)(e)\}_{e \geqslant 0} \subseteq \operatorname{add}_R(X)$. □

Question 4.6. If R is Gorenstein, is $M^* \in \mathcal{FT}(R)$?

Lemma 4.7. *Let $f\colon R \to S$ be a ring homomorphism. Suppose that S is (S_2). Let M be an R-module. Suppose that:*
(a) f is flat; or
(b) M_P is free for every $P = f^{-1}(Q)$ such that $Q \in \operatorname{Spec} S$ and $\operatorname{ht}(Q) = 1$.
If $M \in \mathcal{FT}(R)$, then $M \otimes_R S \in \mathcal{FT}(S)$.

Proof. Let $N = M \otimes_R S$. First, we consider (b) and suppose that $Q \in \operatorname{Spec} S$, $\operatorname{ht}(Q) = 1$ and $M \in \mathcal{FT}(R)$. Let $P = f^{-1}(Q)$. Then $N_Q = M_P \otimes_{R_P} S_Q$, which is free over S_Q by assumption. Next, we consider the modules $F_S^e(N)^{**} = (M \otimes_R S \otimes_S {}^eS)^{**} = (M \otimes_R {}^eS)^{**} = (M \otimes_R {}^eR \otimes_{{}^eR} {}^eS)^{**}$.

$$\begin{array}{ccc} R & \xrightarrow{f} & S \\ {\scriptstyle\varphi}\downarrow & & \downarrow{\scriptstyle\varphi} \\ {}^eR & \xrightarrow{f} & {}^eS \end{array}$$

Consider the maps $R \xrightarrow{\varphi} {}^eR \xrightarrow{f} {}^eS$, where φ is the Frobenius map, and let $R_1 = {}^eR$, $R_2 = {}^eS$. Then Lemma 3.6 shows that $F_S^e(N)^{**} = (M \otimes_R {}^eS)^{**} \cong (F_R^e(M)^{**} \otimes_{{}^eR} {}^eS)^{**}$, so $N(e) \cong (M(e) \otimes_R S)^{**} \subseteq \operatorname{add}_S((X \otimes_R S)^{**})$.
Now suppose that f is flat. Let Q, P and M be as above. Then we have $\operatorname{ht}(Q) = \operatorname{ht}(P) + \dim(S_Q/PS_Q)$, so $\operatorname{ht}(P) \leqslant 1$ and M_P is free, giving (b). □

Lemma 4.8. *Suppose that R is regular. Let M be an R-module. Then M^* is reflexive.*

Proof. Consider the canonical map $f\colon M^* \to M^{***}$. In codimension 1, R is a principal ideal domain, so M^* is free, and f is an isomorphism. Since R is (S_2), so are M^* and M^{***}. By Lemma 3.1, f is an isomorphism. □

Lemma 4.9. *Let $R \to S$ be a flat ring extension. Suppose that M is a reflexive R-module. Then $M \otimes_R S$ is a reflexive S-module.*

Proof. Since $R \to S$ is flat, we have $M \otimes_R S = M^{**} \otimes_R S = (M \otimes_R S)^{**}$. □

Lemma 4.10. *Suppose that R is regular. Consider the following statements:*
(a) $M \in \mathcal{FT}(R)$
(b) M^* is free.
(c) M^{**} is free.
Then (a) $\Rightarrow$ (b) $\Leftrightarrow$ (c). If M is free in codimension 1, then (a) $\Leftrightarrow$ (b) $\Leftrightarrow$ (c).

Proof. (b) $\Rightarrow$ (c): Obvious.
(c) $\Rightarrow$ (b): If M^{**} is free, then so is $M^{***} = (M^*)^{**}$. So M^* is free by Lemma 4.8.
(a) $\Rightarrow$ (c): Suppose that $M \in \mathcal{FT}(R)$. By Proposition 4.5(a) and Lemma 3.1, we may replace M by M^{**} and assume that M is reflexive. So we need to show that M is free. Since R is regular, the ring extension $R \to {}^eR$ is flat. By Lemma 4.9, we have $M(e) = (M \otimes_R {}^eR)^{**} = M \otimes_R {}^eR$ for all $e \geqslant 0$. By Corollary 4.4, we have $M(e') \cong M(e' + e)$ for some $e' \geqslant 0$ and $e > 0$. First, suppose that $e' = 0$, so that $M \cong M(e)$. Consider a minimal free resolution

$$\cdots \to F_1 \xrightarrow{A} F_0 \to M \to 0$$

of M given by the matrix A. Let $I(M) \subseteq \mathfrak{m}$ be the Fitting ideal of M generated by the entries of A and let $q = p^e$. Tensoring with ${}^eR$ gives a free resolution

$$(*) \qquad \cdots \to F_1 \xrightarrow{A^{[q]}} F_0 \to M(e) \to 0,$$

so $I(M(e)) = I(M)^{[q]}$. Since $M \cong M(e)$, we have $I(M) = I(M(e)) = I(M)^{[q]}$, so $I(M) = I(M)^q$. By Nakayama's Lemma, we have $I(M) = 0$. So $A = 0$, and $M^{**} = M$ is free.
In general, the above shows that $M(e)$ is free for some $e \geqslant 0$ by Corollary 3.7. Suppose that $e > 0$. Since $(*)$ is a free resolution of the free module $M(e)$, it is a direct sum of a trivial complex and the resolution $0 \to F \to M(e) \to 0$. So the entries in $A^{[q]}$ are either 1 or 0. Since A has entries in $\mathfrak{m}$ and R is a domain, we must have $A = 0$, so again M is free.
(c) $\Rightarrow$ (a): Suppose that M is free in codimension 1. If M^{**} is free, then $M^{**} \in \mathcal{FT}(R)$, so $M \in \mathcal{FT}(R)$ by Proposition 4.5(a). □

Theorem 4.11. *Let $\varphi\colon R \to S$ be a finite homomorphism of normal domains such that φ is étale in codimension 1 and splits as a map of R-modules. Let N, L be R-modules such that N is (S_2) and that locally, (N, L) is an abundant pair and $N \in \operatorname{add}(L)$. Then $L \in \operatorname{add}_R(\operatorname{CM}(S))$.*

Proof. Since S is of finite F-type over R, $\operatorname{Hom}_R(S, L)$ is C-M by Theorem 3.10, so $\operatorname{Hom}_R(S, L) \in \operatorname{CM}(S)$. Since φ splits, L is a direct summand of $\operatorname{Hom}_R(S, L)$. □

Corollary 4.12. *Let $\varphi\colon R \to S$ be a finite homomorphism of normal domains such that φ is étale in codimension 1 and splits as a map of R-modules and that S is regular (for example, if R is a quotient singularity as in example 5.3). If $M \in \mathcal{FT}(R)$, then $M^* \in \operatorname{add}_R(S)$. If $S^* = \operatorname{Hom}_R(S, R) \cong S$ (again, for example if R is a quotient singularity as in 5.3) then $\mathcal{FT}(R) = \operatorname{add}_R(S)$.*

Proof. Let $N = M \otimes_R S$. If $M \in \mathcal{FT}(R)$, then $N \in \mathcal{FT}(S)$ by Lemma 4.7. Then N^* is free by Lemma 4.10. We have $N^* = \operatorname{Hom}_S(M \otimes_R S, S) = \operatorname{Hom}_R(M, \operatorname{Hom}_S(S, S)) = \operatorname{Hom}_R(M, S)$. Since f splits, $\operatorname{Hom}_R(M, R)$ is a direct summand of $\operatorname{Hom}_R(M, S)$, so $M^* \in \operatorname{add}_R(S)$. If $S^* \cong S$, then it follows that $\mathcal{FT}(R) \subseteq \operatorname{add}_R(S)$, and $S \in \mathcal{FT}(R)$ by Example 4.1. □

Corollary 4.13. *Suppose that k is algebraically closed. Let $S = k[[x_1, \dots, x_d]]$. Let G be a finite subgroup of $GL(d, k)$ that contains no pseudo-reflections such that the order of G is coprime to p. Let $R = S^G$. Then $\mathcal{FT}(R) = \operatorname{add}_R S$.*

Proof. Use Corollary 4.12. □

Theorem 4.14. *Suppose that R is a complete intersection and M is an R-module that is free in codimension 2. Then $M \in \mathcal{FT}(R)$ if and only if M^{**} is free.*

Proof. The proof of "if" is given by (c) ⇒ (a) in Lemma 4.10. For the "only if" part, as in the proof of Lemma 4.10 (a) ⇒ (c), it suffices to assume that M is reflexive and show that M is free. The proof is by induction on d. For $d \geqslant 3$, M is free on $\operatorname{Spec} R \setminus \{\mathfrak{m}\}$ by induction. Let $r = \operatorname{depth} M \geqslant 2$. Suppose that $r < d$. Then by [7, Proposition 4.14], we have $\mathrm{H}^r_{\mathfrak{m}}(F^e_R(M)) \cong \mathrm{H}^r_{\mathfrak{m}}(F^e_R(M)^{**}) = {}^e\mathrm{H}^r_{\mathfrak{m}}(M(e))$ as in Theorem 3.8, so

$$\lim_{e\to\infty} \frac{\lambda_R(\mathrm{H}^r_{\mathfrak{m}}(F^e_R(M)))}{p^{e(d+\alpha(R))}} = \lim_{e\to\infty} \frac{\lambda_R({}^e\mathrm{H}^r_{\mathfrak{m}}(M(e)))}{p^{e(d+\alpha(R))}} = \lim_{e\to\infty} \frac{p^{e\alpha(R)}\lambda_R(\mathrm{H}^r_{\mathfrak{m}}(M(e)))}{p^{e(d+\alpha(R))}} = 0$$

by the assumption that $M \in \mathcal{FT}(R)$. Then [7, Theorem 4.12] gives $\operatorname{pd}_R M < d - r$. By the Auslander-Buchsbaum formula, we have $\operatorname{depth} M > r$, a contradiction. So $r = d$, and again [7, Theorem 4.12] with $k = r - 1$ shows that $\operatorname{pd}_R M = 0$, so M is free. □

Example 4.15. *Let $R = k[[x, y, z]]/(xy - z^2)$ and M be the ideal (x, z). Then $[M]$ has order 2 in $\mathrm{Cl}(R)$ and so $M \in \mathcal{FT}(R)$ by Example 4.1, but M^{**} is not free, so the condition that M is locally free in codimension 2 is necessary in Theorem 4.14.*

5. F-abundant pairs and modules

In this section we give many examples of F-abundant pairs and modules.

Example 5.1. *(a) If $\operatorname{sdim} R \geqslant 1$, in particular if R is strongly F-regular of dimension $\geqslant 1$, then (R, R) is an abundant pair.*
(b) [14, Proposition 2.3] *shows that $\alpha(R_P) = \alpha(R) + \dim(R/P)$. Let N, L be as in Definition 1.1 and $P \in \operatorname{Spec} R$. If $\liminf_{e\to\infty} p^{e(\alpha(R)+\dim(R/P))}/b_e = 0$, then (N_P, L_P) is an abundant pair.*
(c) F-contributors for modules of finite F-representation type, as in [22, Section 2].

Example 5.2. *(*[7, Example 6.1]*) Let k be an algebraically closed field of characteristic $p > 2$. Consider the hypersurface $R = k[[x, y, u, v]]/(xy - uv)$. Then every MCM R-module is F-abundant.*

Example 5.3. *Let k be an algebraically closed field of characteristic $p > 0$ and V be a k-vector space of dimension d. Let S be the symmetric algebra of V. Let G be a finite subgroup of $GL(V)$ without pseudo-reflections such that $|G|$ is coprime to p. Let $R = S^G$ be the ring of invariants. Let $V_0, \cdots, V_n$ be a complete set of irreducible representations of G over k.*

Let $M_i = (S \otimes_k V_i)^G$. *It is classical that* $\operatorname{add}_R(S) = \{M_0, M_1, \dots, M_n\}$. *Also, by the main results of* [10]*:*

(1) S*, and hence all of* $M_0, M_1, \dots, M_n$*, are modules of finite* F*-type. Note that* $\operatorname{rank} M_i = \dim_k V_i$*, so we have many examples of modules of finite* F*-type which are not ideals.*
(2) (M_i, M_j) *is an* F*-abundant pair for all* $0 \le i, j \le n$.

Corollary 5.4. *Let* R, N *be as in Theorem 3.8. Suppose that* $\liminf_{e\to\infty} p^{e(\alpha(R)+d-3)}/b_e = 0$ *as in Example 5.1 with* $N = L$ *(for example, when* $d = 3$ *and* (N, N) *is an abundant pair). Then* N *is MCM. In particular, if* R *is regular, then* N *is free.*

Proof. Theorem 3.10 with $M = R$ and $e = 0$ shows that N is MCM. □

Example 5.5. *Let* k *be a perfect field,* $R = k[[x, y, z]]$ *and* M *be the ideal* (x, y)*. Then* (M, M) *is an abundant pair, but* M *is not* (S_2)*. So the assumption in Corollary 5.4 for* N *to be* (S_2) *cannot be weakened.*

Proof. Let $C = k[[z]]$. Consider the exact sequence

$$0 \to M \to R \to C \to 0$$

Let $q = p^e$. Since ${}^e-$ is an exact functor, we have

$$\begin{array}{ccccccccc} 0 & \longrightarrow & {}^eM & \longrightarrow & {}^eR & \longrightarrow & {}^eC & \longrightarrow & 0 \\ & & & & \| & & \| & & \\ & & & & R^{\oplus q^3} & \longrightarrow & C^{\oplus q} & \longrightarrow & 0 \end{array}$$

It follows that ${}^eM$ has exactly q copies of M, so (M, M) is an abundant pair. Since $1 = \operatorname{depth} M_M < \dim M_M = 2$, M is not (S_2). □

Example 5.6. *Let* k *be a perfect field,* $R = k[[x_1, x_2, \dots, x_{d-3}, u, v, w]]$, $C = R/(u, v, w)$ *and* $M = \Omega^2 C$*, the second syzygy of* C*. Then* M *is* (S_2) *and* (M, M) *is an abundant pair. So the assumptions in Corollary 5.4 cannot be weakened.*

Proof. Let $q = p^e$. Then as in Example 5.5, we have

$$\begin{array}{ccccccccccc} 0 & \longrightarrow & {}^eM & \longrightarrow & {}^e(R^3) & \longrightarrow & {}^eR & \longrightarrow & {}^eC & \longrightarrow & 0 \\ & & & & \| & & \| & & \| & & \\ & & & & R^{\oplus 3q^d} & \longrightarrow & R^{\oplus q^d} & \longrightarrow & C^{\oplus q^{d-3}} & \longrightarrow & 0 \end{array}$$

It follows that ${}^eM$ has exactly q^{d-3} copies of M. Then as in Corollary 5.4, we have $\liminf_{e\to\infty} q^{(\alpha(R)+d-3)}/b_e = q^{d-3}/q^{d-3} = 1$. Since M is a second syzygy, it is (S_2). Since $\operatorname{depth} C = d - 3$, we have $\operatorname{depth} M = d - 1 \ne d$, so M is not free. □

6. Geometric applications

Discussion 6.0. Let $(A, \mathfrak{m})$ be a standard graded ring that is (S_2) of dimension at least 2. We first fix some notation and record some results from [9, 16]. Let $R = A_{\mathfrak{m}}$. Let $X = \mathbf{Proj}(A)$, $Y = \mathbf{Spec}(R) \setminus \{\mathfrak{m}_{\mathfrak{m}}\}$ and $Z = \mathbf{Spec}(R)$. Let $\iota\colon Y \to Z$ be the inclusion morphism. There

is also an affine surjective morphism $\pi\colon Y \to X$ ([16, Proposition I.5]). Corollary I.6 of [16] states that for every A-graded module M we have sheaf cohomology isomorphisms

$$\oplus_{d\in\mathbb{Z}} H^i(X, \widetilde{M}(d)) \xrightarrow{\simeq} H^i(Y, \widetilde{M_{\mathfrak{m}}|_Y})$$

for $i \geqslant 0$. For $1 \leqslant i \leqslant d-1$, we have $H^i(Y, \widetilde{M_{\mathfrak{m}}|_Y}) \cong H^{i+1}_{\mathfrak{m}}(M_{\mathfrak{m}})$.

Let $V(\cdot)$ denote the category of vector bundles over a scheme. Let Γ be the global section functor. If $\mathscr{G} \in V(Y)$, then $\iota_*\mathscr{G}$ is a coherent $\mathscr{O}_Z$-module. Let $\psi = \Gamma \circ \pi^*$. Then we have maps

$$V(X) \xrightarrow{\pi^*} V(Y) \xrightarrow{\Gamma} \{M \in \operatorname{Ref}(R) \mid M \text{ is locally free on } Y\}$$

By [9, Theorem 1.3], $\operatorname{depth}\Gamma(\mathscr{G}) \geqslant 2$. Γ satisfies the property $\Gamma(\mathscr{G}_1 \otimes \mathscr{G}_2) = (\Gamma(\mathscr{G}_1) \otimes_R \Gamma(\mathscr{G}_2))^{**}$. Let $\mathscr{F}_i \in V(X)$, $i = 1, 2$, be indecomposable such that $\psi(\mathscr{F}_i)$ are isomorphic up to free summand. Then by [9, Proposition 9.5], $\mathscr{F}_1 \cong \mathscr{F}_2(m) = \mathscr{F}_2 \otimes \mathscr{O}_X(m)$ for some m.

Definition 6.1. ([4, Introduction]) Let X be a smooth projective variety defined over a field k of characteristic $p > 0$. Let $\varphi\colon X \to X$ be the absolute Frobenius morphism. Then a vector bundle $\mathscr{F} \in V(X)$ is (e, f)-Frobenius periodic, or (e, f)-F periodic in short, if there are $e < f$ such that $(\varphi^e)^*(\mathscr{F}) \cong (\varphi^f)^*(\mathscr{F})$.

Lemma 6.2. *Suppose that $\mathscr{F} \in V(X)$ is (e, f)-F periodic. Let X, Y be as in 6.0. Let $M = \psi(\mathscr{F})$. Then $\mathscr{F}$ is (e_1, e_2)-F periodic iff $M(e_1) \cong M(e_2)$.*

Proof. For "only if", we have $\psi\big((\varphi^e)^*(\mathscr{F})\big) = \psi(\mathscr{F} \otimes (\varphi^e)^*(X)) = \big(\psi(\mathscr{F}) \otimes (\varphi^e)^*(X)\big)^{**} = M(e)$ as in 6.0. The "if" part comes from sheafification. □

Next, we discuss a generalization of one of the results in [18, Theorem 3.1]. For that we need to recall some notation. Let R be a F-finite normal domain and let D be a $\mathbb{Q}$-divisor.

In the next result we are able to remove the condition that the characteristic p is coprime to r as in [18, Theorem 3.1]. We follow the same trick as in [18], with a crucial difference suggested by our approach in this paper: the reflexive module representing a torsion element in the class group has finite F-type.

Theorem 6.3. *Let R be an F-finite normal domain with perfect residue field and $X = \operatorname{Spec} R$. Let Δ be a $\mathbb{Q}$-divisor on X such that the pair (X, Δ) is strongly F-regular. Let D be an integral divisor such that $rD \sim r\Delta'$ for some integer $r > 0$ and $0 \leq \Delta' \leq \Delta$. Then $\mathscr{O}_X(-D)$ is Cohen-Macaulay.*

Proof. Since we also have that the pair (X, Δ') is strongly F-regular, one may assume $\Delta' = \Delta$. Now, this assumption implies that there is a decomposition of R-modules ($q = p^e$) ([5, Lemma 3.5]):

$$F^e_*\mathscr{O}_X((q-1)\Delta) = \mathscr{O}_X^{n_e} \oplus N_e$$

such that $\liminf_{e\to\infty} \frac{n_e}{q^d} > 0$. Twisting by $\mathscr{O}_X(-D)$, reflexifying, we get a decomposition:

$$F^e_*\mathscr{O}_X((q-1)(\Delta - D) - D) = \mathscr{O}_X(-D)^{n_e} \oplus N'_e$$

The key point now is that as $r(\Delta - D) \sim 0$, there are only finitely many isomorphism classes of the modules $\mathscr{O}_X((q-1)(\Delta - D) - D)$. Let M be the direct sum of all these modules and $I = \mathscr{O}_X(-D)$, what we have is precisely:

$${}^eM \cong I^{n_e} \oplus P_e$$

with $\liminf_{e\to\infty} \frac{n_e}{q^d} > 0$. Lemma 2.7 now forces I to be Cohen-Macaulay. □

7. Acknowledgments

The first author would like to thank Kevin Tucker and Karl Schwede for patiently explaining to him many basic facts on the subjects this note.

References

1. *Twenty-Four Hours of Local Cohomology*, S. Iyengar, G. J. Leuschke, A. Leykin, C. Miller, E. Miller, A. K. Singh, U. Walther, American Mathematical Society, Providence, R.I., c2007.
2. *Local Cohomology*, M. P. Brodmann, R. Y. Sharp, Cambridge Studies in Advanced Mathematics **136**, Cambridge University Press, 2013.
3. I. M. Aberbach, F. Enescu, *The Structure of F-Pure Rings*, Math. Z. **250** (2005), no. 4, 791–806.
4. H. Brenner, A. Kaid, *An Explicit Example of Frobenius Periodicity*, J. Pure Appl. Algebra **217** (2013), no. 8, 1412–1420.
5. M. Blickle, K. Schwede, K. Tucker, *F-signature of Pairs and the Asymptotic Behavior of Frobenius Splittings*, Adv. in Math. **231** (2012), 3232–3258.
6. N. R. Baeth, R. Wiegand, *Factorization Theory and Decompositions of Modules*, American Mathematical Monthly **120**, no. 1 (January 2013), 3–34.
7. H. Dao, I. Smirnov, *On Generalized Hilbert-Kunz Function and Multiplicity*, arXiv:1305.1833.
8. K. R. Fuller, W. A. Shutters, *Projective Modules over Non-commutative Semilocal Rings*, Tohoku Math. J. **27** (1975), no. 3, 303–311.
9. G. Horrocks, *Vector Bundles on the Punctured Spectrum of a Local Ring*, Proc. London Math. Soc. (3) **14** (1964), 689–713.
10. M. Hashimoto, Y. Nakajima, *Generalized F-signature of invariant subrings*, preprint `http://front.math.ucdavis.edu/1311.5963`.
11. J. Horiuchi, L. E. Miller, K. Shimomoto, Appendix by K. Schwede, A. K. Singh, *Deformation of F-Injectivity and Local Cohomology*, arXiv:1210.3589v3.
12. N. M. Katz, *p-adic Properties of Modular Schemes and Modular Forms*, Modular Functions of One Variable, III (Proc. Internat. Summer School, Univ. Antwerp, Antwerp, 1972), 69–190. Lecture Notes in Mathematics, **350**, Springer, Berlin, 1973.
13. J. Kollár, *A Local Version of the Kawamata-Viehweg Vanishing Theorem*, Pure Appl. Math. Q. **7** (2011), no. 4, Special Issue: In memory of Eckart Viehweg, 1477–1494.
14. E. Kunz, *On Noetherian Rings of Characteristic p*, American J. of Math. **98** (Winter, 1976), no. 4, 999–1013.
15. V. B. Mehta, V. Srinivas, *Varieties in Positive Characteristic with Trivial Tangent Bundle*, Compositio Mathematica **64.2** (1987), 191–212.
16. M. Majidi-Zolbanin, *Splitting of Vector Bundles on Punctured Spectrum of Regular Local Rings*, Doctoral dissertation, the Graduate Center of the City University of New York (2005).
17. Y. Nakajima, *Dual F-Signature of Special Cohen-Macaulay Modules over Cyclic Quotient Surface Singularities*, arXiv:1311.5967v1.
18. Z. Patakfalvi, K. Schwede, *Depth of F-singularities and Base Change of Relative Canonical Sheaves*, J. Inst. Math. Jussieu **13** (2014), no. 1, 43–63.
19. K. Smith, *Globally F-regular Varieties: Applications to Vanishing Theorems for Quotients of Fano Varieties*, Michigan Math. J. **48** (2000), no. 1, 553–572.
20. K. Tucker, *F-Signature Exists*, Inventiones mathematicae **190** (2012), 743–765.
21. K. Watanabe, *F-regular and F-pure normal graded rings*, Journal of Pure and Applied Algebra **71** (1991), 341–350.
22. Y. Yao, *Modules with Finite F-Representation Type*, J. London Math. Soc. (2) **72** (2005), no. 1, 53–72.

Department of Mathematics, University of Kansas, Lawrence, KS 66045-7523 USA
E-mail address: hdao@ku.edu

Department of Mathematics, University of Kansas, Lawrence, KS 66045-7523 USA
E-mail address: tonyse@ku.edu